

REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 10/28/94	3. REPORT TYPE AND DATES COVERED Final, 10/1/91 - 9/30/94	
4. TITLE AND SUBTITLE Coupling of Brain Muscarinic Receptors to Second Messengers			5. FUNDING NUMBERS DAAL03-91-G-0337	
6. AUTHOR(S) Esam E. El-Fakahany, Ph.D., Div. of Neuroscience Research in Psychiatry, University of MN Med. School, Box 392 UMHC, Minneapolis, MN 55455			8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Minnesota, Office of Research & Technology Transfer Administration, 1100 Washington Ave. S.E., Suite 201, Minneapolis, MN 55415			9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709-2211	
11. SUPPLEMENTARY NOTES The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.			12b. DISTRIBUTION CODE ARO 29594.1-L5	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			13. ABSTRACT (Maximum 200 words) Stimulating muscarinic acetylcholine receptors results in activation of various signal transduction pathways. The goal of the research was to investigate the selectivity of coupling of the currently known 5 subtypes of muscarinic receptors to these different second messenger pathways. Results indicate that m1 m3 and m5 muscarinic receptors are coupled to enhanced phosphoinositide hydrolysis, while m2 and m4 receptors are coupled to a decrease in cyclic AMP formation. Results also demonstrate efficient coupling of m5 muscarinic receptors to generation of nitric oxide.	
14. SUBJECT TERMS Receptors - Muscarinic acetylcholine - second messengers - signal transduction.			15. NUMBER OF PAGES	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED			16. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED			19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	
20. LIMITATION OF ABSTRACT UL			21. LIMITATION OF ABSTRACT UL	

FINAL REPORT

29594-LS

"Coupling of Brain Muscarinic Receptors to Second Messengers"

Esam E. El-Fakahany, Ph.D.

STATEMENT OF THE PROBLEM STUDIED:

Muscarinic acetylcholine receptors play very important roles in relation to regulation of brain functions. Mammalian brain contains five different subtypes of muscarinic cholinergic receptors which respond to the neurotransmitter acetylcholine. The main goal of the research project was to investigate the selectivity of coupling of these receptor subtypes to various second messenger signaling mechanisms.

SUMMARY OF THE MOST IMPORTANT RESULTS:

The selectivity of coupling of different subtypes of muscarinic receptors to various second messenger signaling pathways was studied in mammalian cell lines which are individually transfected with the genes which encode for each of the five subtypes of muscarinic acetylcholine receptors. Activation of m1, m3 or m5 muscarinic receptors resulted in a significant enhancement of the hydrolysis of phosphoinositides. On the other hand, activation of m2 and m4 muscarinic receptors resulted in inhibition of cyclic AMP formation. Some muscarinic receptor agonists are able to discriminate among various subtypes of the receptor by showing functional selectivity. Our studies demonstrated the importance of equalizing the number of expressed receptors for studying such agonist selectivity.

We also studied the selectivity of coupling of various subtypes of muscarinic receptors to another very important second messenger signaling mechanism, namely the generation and release of nitric oxide. Nitric oxide plays important roles in learning and memory. It had also been shown to be involved in neurotoxicity. Our studies demonstrated that m5 muscarinic receptors which are efficiently coupled to enhanced generation of nitric oxide. These receptors are localized in basal ganglia. Their efficient coupling to generation of nitric oxide might be important for cholinergic enhancement of dopamine release.

PUBLICATIONS:

1. Kiefer-Day, J. and E.E. El-Fakahany: Muscarinic Receptor Function and Acetylcholinesterase Activity After Chronic Administration of Tacrine to Mice at Therapeutic Drug Concentrations. *Pharmacology*, 44, 71-80, 1992.
2. Lee, N.H., J. Hu and E.E. El-Fakahany: Modulation by Certain Conserved Aspartate Residues of the Allosteric Interaction of Gallamine at the m1 Muscarinic Receptor. *Journal of Pharmacology and Experimental Therapeutics*, 262, 312-316, 1992 .
3. Meng, Y., J. Hu and E.E. El-Fakahany: p-Fluoro-hexahydro-siladifenidol Exhibits Poor Selectivity between M₃ and M₁ Muscarinic Receptors. *Membrane Biochemistry*, 9, 293-300, 1992.
4. Hu, J., S.H. Wang and E.E. El-Fakahany: Role of Conserved Aspartate Residue 71 in Down-Regulation of M1 Muscarinic Receptors. *Pharmacology Communications*, 1, 219-225, 1992.
5. Zhu, S.Z., S.H. Wang, D.C. Mash and E.E. El-Fakahany: Comparison of the Concentration of Messenger RNA Encoding Four Muscarinic Receptor Subtypes in Control and Alzheimer Brains. *Molecular Brain Research*, 16, 64-70, 1992.
6. Hu, J., S.H. Wang, C. Forray and E.E. El-Fakahany: Complex Allosteric Modulation of Cardiac Muscarinic Receptors by Protamine: a Potential Model for Putative Endogenous Ligands. *Molecular Pharmacology*, 42, 311-324, 1992.
7. Mohamed, A.S., C. Forray and E.E. El-Fakahany: Lack of Intrinsic Activity and Significant Subtype Selectivity of SR 95639A at Muscarinic Receptors. *European Journal of Pharmacology (Molecular Pharmacology Section)*, 227, 181-187, 1992.
8. Wang, S.Z., S.Z. Zhu and E.E. El-Fakahany: Comparison of the Level of mRNA Encoding M1 and M2 Muscarinic Receptors in Brains of Young and Aged Rats. *Neuroscience Letters*, 145, 149-152, 1992.
9. Hu, J. and E.E. El-Fakahany: The Calmodulin Antagonist Calmidazolium Stimulates Release of Nitric Oxide in Neuroblastoma N1E-115 Cells. *NeuroReport*, 4, 198-200, 1993.

10. Hu, J. and E.E. El-Fakahany: Role of Intercellular and Intracellular Communication by Nitric Oxide in Coupling of Muscarinic Receptors to Activation of Guanylate Cyclase in Neuronal Cells. *Journal of Neurochemistry*, 61, 578-585, 1993.
11. Kiefer-Day, J., E.A.M. Abdallah, C. Forray, N.H. Lee, O.N. Kim and E.E. El-Fakahany: Effects of Tacrine on Brain Muscarinic Receptor-Mediated Second Messenger Signals. *Pharmacology*, 47, 98-110, 1993.
12. Wang, S.Z. and E.E. El-Fakahany: Application of Transfected Cell Lines in Studies of Functional Receptor Subtype Selectivity of Muscarinic Agonists. *Journal of Pharmacology and Experimental Therapeutics*, 266, 237-243, 1993.
13. Hu, J. and E.E. El-Fakahany: β -Amyloid 25-35 Activates Nitric Oxide Synthase in a Neuronal Clone. *NeuroReport*, 4, 670-672, 1993.
14. Hu, J. and E.E. El-Fakahany: Allosteric Interaction of Dynorphin and Myelin Basic Protein with Muscarinic Receptors. *Pharmacology*, 47, 351-359, 1994.
15. Hu, J., J.H. Lee and E.E. El-Fakahany: Inhibition of Neuronal Nitric Oxide Synthase by Antipsychotic Drugs. *Psychopharmacology*, 114, 161-166, 1994.
16. Wang, S.Z., S.Z. Zhu and E.E. El-Fakahany: Efficient Coupling of m5 Muscarinic Acetylcholine Receptors to Activation of Nitric Oxide Synthase. *Journal of Pharmacology and Experimental Therapeutics*, 268, 552-557, 1994.
17. Zhu, S.Z., S.Z. Wang, J. Hu and E.E. El-Fakahany: An Arginine Residue Conserved in Most G-Protein-Coupled Receptors is Essential for the Function of m1 Muscarinic Receptors. *Molecular Pharmacology*, 45, 517-523, 1994.
18. Iadecola, C., X. Xu, F. Zhang, J. Hu and E.E. El-Fakahany: Prolonged Inhibition of Brain Nitric Oxide Synthase by Short-Term Systemic Administration of Nitro-L-arginine Methyl Ester. *Neurochemical Research*, 19, 501-505, 1994.
19. Hu, J., M.I. Mahmoud and E.E. El-Fakahany: Polyamines Inhibit Nitric Oxide Synthase in Rat Cerebellum. *Neuroscience Letters*, 175, 41-45, 1994.
20. Hu, J. and E.E. El-Fakahany: An Artifact Associated with Using Trypan Blue Exclusion to Measure Effects of Amyloid β on Neuron Viability. *Life Sciences*, 55, 1009-1016, 1994.

LIST OF PARTICIPATING SCIENTIFIC PERSONNEL:

Sheng Zu Zhu

No degrees granted while employed on the project.

REPORT OF INVENTIONS: None.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	